

GROTHENDIECK LOCAL DUALITY AND COHOMOLOGICAL HASSE PRINCIPLE FOR 2-DIMENSIONAL COMPLETE LOCAL RING

BELGACEM DRAOUIL

ABSTRACT. We prove a local duality for some schemes associated to a 2-dimensional complete local ring whose residue field is an n -dimensional local field in the sense of Kato-Parshin. Our results generalize the Saito works in the case $n = 0$ and are applied to study the Bloch-Ogus complex for such rings in various cases.

1. INTRODUCTION

Let A be a 2-dimensional complete local ring with finite residue field. The Bloch-Ogus complex associated to A has been studied by Saito in [12]. In this prospect, he calculated the homologies of this complex and obtained (for any integer $n \geq 1$) the following exact sequence

$$(1.1) \quad 0 \longrightarrow (\mathbb{Z}/n)^r \longrightarrow H^3(K, \mathbb{Z}/n(2)) \longrightarrow \bigoplus_{v \in P} H^2(k(v), \mathbb{Z}/n(1)) \longrightarrow \mathbb{Z}/n \longrightarrow 0,$$

where P denotes the set of height one prime ideals of A , K is the fractional field of A , $k(v)$ is the residue field at $v \in P$, and $r = r(A)$ is an integer depending on the degeneracy of $\text{Spec} A$. This result is based upon the isomorphism ([12], lemma 5.4):

$$(1.2) \quad H^4(X, \mathbb{Z}/n(2)) \simeq \mathbb{Z}/n,$$

where $X = \text{Spec} A \setminus \{x\}$; and x is the unique maximal ideal of A . A decade later, Matsumi in [8] generalised the result by Saito to 3-dimensional complete regular local ring of positive characteristic. Indeed, he proved the exactness of the complex

$$(1.3) \quad \begin{aligned} 0 \longrightarrow H^4(K, \mathbb{Z}/\ell(3)) &\longrightarrow \bigoplus_{v \in (\text{Spec} A)_2} H^3(k(v), \mathbb{Z}/\ell(2)) \\ &\longrightarrow \bigoplus_{v \in (\text{Spec} A)_1} H^2(k(v), \mathbb{Z}/\ell(1)) \longrightarrow \mathbb{Z}/\ell \longrightarrow 0 \end{aligned}$$

for all ℓ prime to $\text{char}(A)$, where $(\text{Spec} A)_i$ indicates the set of all points in $\text{Spec} A$ of dimension i . Besides, if the ring A is not regular, then the map

$$H^4(K, \mathbb{Z}/\ell(3)) \xrightarrow{\Psi_K} \bigoplus_{v \in (\text{Spec} A)_2} H^3(k(v), \mathbb{Z}/\ell(2))$$

is non-injective.

We proved in ([3], th3) that $\text{Ker} \Psi_K$ contains a sub-group of type $(\mathbb{Z}/\ell)^{r'_1(A)}$, where $r'_1(A)$ is calculated as the $\mathbb{Z}$ -rank of the graph of the exceptional fiber of a resolution of $\text{Spec} A$. The main tools used in this direction are the isomorphism

$$(1.4) \quad H^6(X, \mathbb{Z}/\ell(3)) \simeq \mathbb{Z}/\ell$$

Date: November 2006.

1991 Mathematics Subject Classification. 11G20, 11G45, 14H30, 14C35, 19F05.

Key words and phrases. Hasse principle, Purity, local duality.

and the perfect pairing

$$(1.5) \quad H^i(X, \mathbb{Z}/\ell) \times H^{6-i}(X, \mathbb{Z}/\ell(3)) \longrightarrow H^6(X, \mathbb{Z}/\ell(3)) \simeq \mathbb{Z}/\ell$$

for all $i \geq 1$ ([3], Section 0,D2).

In order to generalize the previous results, we consider a 2-dimensional complete local ring A whose residue field k is n -dimensional local field in the sense of Kato-Parshin. To be more precise, let $X = \text{Spec} A \setminus \{x\}$, where x is the unique maximal ideal of A . Then the main result of this paper is the following

Theorem (theorem 3.1)

There exist an isomorphism

$$(1.6) \quad H^{4+n}(X, \mathbb{Z}/\ell(2+n)) \simeq \mathbb{Z}/\ell$$

and a perfect pairing

$$(1.7) \quad H^1(X, \mathbb{Z}/\ell) \times H^{3+n}(X, \mathbb{Z}/\ell(2+n)) \longrightarrow H^{4+n}(X, \mathbb{Z}/\ell(2+n)) \simeq \mathbb{Z}/\ell$$

for all ℓ prime to $\text{char}(A)$.

We apply this result to calculate the homologies of the Bloch-Ogus complex associated to A . Indeed, let $\pi_1^{c.s.}(X)$ be the quotient group of $\pi_1^{ab}(X)$ which classifies abelian c.s coverings of X (see definition 4.1 below). We then prove the following

Theorem (theorem 4.2)

Let A be a 2-dimensional complete normal local ring of positive characteristic whose residue field is n -dimensional local field. Then the exact sequence

$$(1.8) \quad \begin{aligned} 0 \longrightarrow \pi_1^{c.s.}(X)/\ell &\longrightarrow H^{3+n}(K, \mathbb{Z}/\ell(2+n)) \\ &\longrightarrow \bigoplus_{v \in P} H^{2+n}(k(v), \mathbb{Z}/\ell(1+n)) \longrightarrow \mathbb{Z}/\ell \longrightarrow 0 \end{aligned}$$

holds.

Furthermore, if A is assumed to be regular then, as in the case of $n = 0$ considered by Saito [12], we prove the vanishing of the group $\pi_1^{c.s.}(X)$ using the recent paper [11] by Panin.

To prove these results, we rely heavily on the Grothendieck duality theorem for strict local rings (section 3) as well as the purity theorem of Fujiwara-Gabber, which we recall next.

In ([4], sentences just below Corollary 7.1.7), Fujiwara confirmed that the absolute cohomological purity in equicharacteristic is true. In other words, we get the following.

Theorem of Fujiwara-Gabber

Let T be an equicharacteristic Noetherian excellent regular scheme and Z be a regular closed subscheme of codimension c . Then for an arbitrary natural number ℓ prime to $\text{char}(T)$, the following canonical isomorphism

$$(1.9) \quad H_Z^i(T, \mathbb{Z}/\ell(j)) \simeq H^{i-2c}(Z, \mathbb{Z}/\ell(j-c))$$

holds.

Finally, we complete the partial duality (1.7) in the case $n = 1$. So, we obtain the following.

Theorem (theorem 5.1)

Let A be a 2-dimensional normal complete local ring whose residue field is one-dimensional local field. Then, for very ℓ prime to $\text{char}(A)$, the isomorphism

$$(1.10) \quad H^5(X, \mathbb{Z}/\ell(3)) \simeq \mathbb{Z}/\ell$$

and the perfect pairing

$$(1.11) \quad H^i(X, \mathbb{Z}/\ell) \times H^{5-i}(X, \mathbb{Z}/\ell(3)) \longrightarrow H^5(X, \mathbb{Z}/\ell(3)) \simeq \mathbb{Z}/\ell$$

hold for all $i \in \{0, \dots, 5\}$

Our paper is organised as follows. Section 2 devoted to some notations. Section 3 contains the main theorem of this work concerning the duality of the scheme $X = \text{Spec} A \setminus \{x\}$, where A is a 2-dimensional complete local ring whose residue field is n -dimensional local field and x is the unique maximal ideal of A . In section 4, we study the Bloch-Ogus complex associated to A . In section 5, we investigate the particular case $n = 1$.

2. NOTATIONS

For an abelian group M and a positive integer n , we denote by M/n the cokernel of the map $M \xrightarrow{n} M$. For a scheme Z , and a sheaf $\mathcal{F}$ over the étale site of Z , $H^i(Z, \mathcal{F})$ denotes the i -th étale cohomology group. For a positive integer ℓ invertible on Z , $\mathbb{Z}/\ell(1)$ denotes the sheaf of ℓ -th root of unity and for an integer i , we denote $\mathbb{Z}/\ell(i) = (\mathbb{Z}/\ell(1))^{\otimes i}$.

A local field k is said to be *n -dimensional local* if there exists the following sequence of fields k_i ($1 \leq i \leq n$) such that

- (i) each k_i is a complete discrete valuation field having k_{i-1} as the residue field of the valuation ring O_{k_i} of k_i , and
- (ii) k_0 is a finite field.

For such a field, and for ℓ prime to $\text{Char}(k)$, the well-known isomorphism

$$(2.1) \quad H^{n+1}(k, \mathbb{Z}/\ell(n)) \simeq \mathbb{Z}/\ell$$

and for each $i \in \{0, \dots, n+1\}$ a perfect duality

$$(2.2) \quad H^i(k, \mathbb{Z}/\ell(j)) \times H^{n+1-i}(k, \mathbb{Z}/\ell(n-j)) \longrightarrow H^{n+1}(k, \mathbb{Z}/\ell(n)) \simeq \mathbb{Z}/\ell$$

hold.

For a field L , $K_i(L)$ is the i -th Milnor group. It coincides with the i -th Quillen group for $i \leq 2$. For ℓ prime to $\text{char } L$, there is a Galois symbol

$$(2.3) \quad h_{\ell, L}^i : K_i L / \ell \longrightarrow H^i(L, \mathbb{Z}/\ell(i))$$

which is an isomorphism for $i = 0, 1, 2$ ($i = 2$ is Merkur'jev-Suslin). In this context, we recall the Kato conjecture ([7], Conjecture 1, Section 1):

Kato Conjecture

For any field L and any ℓ prime to $\text{char } L$, the map $h_{\ell, L}^i$ is bijective.

3. LOCAL DUALITY

We start this section by a description of the Grothendieck local duality. Let B denote a d -dimensional normal complete local ring with maximal ideal x' . By Cohen structure theorem ([10], 31.1), B is a quotient of a regular local ring. Hence $\text{Spec } B$ admits a dualizing complex. Now, assume in the first step that the residue field of B is separably closed (B is strictly local). Then, for $X' = \text{Spec } B \setminus \{x'\}$ and for any ℓ prime to $\text{char}(A)$, there is a Poincaré duality theory ([15], Exposé I, Remarque 4.7.17). Namely, there is a trace isomorphism

$$(3.1) \quad H^{2d-1}(X', \mathbb{Z}/\ell(d)) \xrightarrow{\sim} \mathbb{Z}/\ell$$

and a perfect pairing

$$(3.2) \quad H^i(X', \mathbb{Z}/\ell) \times H^{2d-1-i}(X', \mathbb{Z}/\ell(d)) \longrightarrow H^{2d-1}(X', \mathbb{Z}/\ell(d)) \simeq \mathbb{Z}/\ell$$

for all $i \in \{0, \dots, 2d-1\}$.

Assume at this point that the residue field k of B is arbitrary. Let k_s be a separable closure of k . The strict henselization B^{sh} of B (with respect to the separably closed extension k_s of k) at the unique maximal ideal x of B is a strictly local ring. It coincides with the integral closure of B in the maximal unramified extension L^{ur} of the fraction field L of B . Let x' be the maximal ideal of B^{sh} and let $X' = \text{Spec} B^{sh} \setminus \{x'\}$. So, the Galois group of X' over X is $\text{Gal}(L^{ur}/L)$ which is isomorphic to $\text{Gal}(k_s/k)$. Then for any integer $j \geq 0$, we get the Hochschild-Serre spectral sequence ([9], Remark 2.21)

$$(3.3) \quad E_2^{p,q} = H^p(k, H^q(X', \mathbb{Z}/\ell(j))) \implies H^{p+q}(X, \mathbb{Z}/\ell(j))$$

Let A denote a 2-dimensional normal complete local ring whose residue field is an n -dimensional local field. Let x be the unique maximal ideal of A . Then by normality A admits at most one singularity at x in such a way that the scheme $X = \text{Spec} A \setminus \{x\}$ becomes a regular scheme.

In what follows, we put

K : the fractional field of A ,

k : the residue field of K

P : the set of height one prime ideals of A .

For each $v \in P$ we denote by K_v the completion of K at v and by $k(v)$ the residue field of K_v .

Let $X = \text{Spec} A \setminus \{x\}$ as above. Generalizing (1.2), (1.4), and (1.5), we get the following.

Theorem 3.1

For all ℓ prime to $\text{char}(A)$, the isomorphism

$$(3.4) \quad H^{4+n}(X, \mathbb{Z}/\ell(2+n)) \simeq \mathbb{Z}/\ell$$

and the perfect pairing

$$(3.5) \quad H^1(X, \mathbb{Z}/\ell) \times H^{3+n}(X, \mathbb{Z}/\ell(2+n)) \longrightarrow H^{4+n}(X, \mathbb{Z}/\ell(2+n)) \simeq \mathbb{Z}/\ell$$

occur. Furthermore, this duality is compatible with duality (2.2) in the sense that the commutative diagram

$$(3.6) \quad \begin{array}{ccccccc} H^1(X, \mathbb{Z}/\ell) & \times & H^{3+n}(X, \mathbb{Z}/\ell(2+n)) & \longrightarrow & H^{4+n}(X, \mathbb{Z}/\ell(2+n)) & \xrightarrow{\sim} & \mathbb{Z}/\ell \\ \downarrow i^* & & \uparrow i_* & & \uparrow i_* & & \parallel \\ H^1(k(v), \mathbb{Z}/\ell) & \times & H^{n+1}(k(v), \mathbb{Z}/\ell(n+1)) & \longrightarrow & H^{n+2}(k(v), \mathbb{Z}/\ell(n+1)) & \xrightarrow{\sim} & \mathbb{Z}/\ell \end{array}$$

holds, where i^* is the map on H^i induced from the map $v \longrightarrow X$ and i_* is the Gysin map.

Proof. The proof is slightly different from the proof of Theorem 1 in [2]. Let k_s be a separable closure of k . We consider the strict henselization A^{sh} of A (with respect to the separably closed extension k_s of k) at the unique maximal ideal x of A . Then, we denote x' the unique maximal ideal of A^{sh} , $X' = \text{Spec} A^{sh} \setminus \{x'\}$ and we use the spectral sequence (3.3). As k is n -dimensional local field, we have $H^{n+2}(k, M) = 0$ for any torsion module M and as X' is of cohomological dimension $2d-1$ ([14], the last paragraph of Introduction), we obtain

$$\begin{aligned} H^{4+n}(X, \mathbb{Z}/\ell(2+n)) &\simeq H^{n+1}(k, H^3(X', \mathbb{Z}/\ell(2+n))) \\ &\simeq H^{n+1}(k, \mathbb{Z}/\ell(n)) \quad \text{by (3.1)} \\ &\simeq \mathbb{Z}/\ell \quad \text{by (2.1)} \end{aligned}$$

We prove now the duality (3.5). The filtration of the group $H^{3+n}(X, \mathbb{Z}/\ell(2+n))$ is

$$H^{3+n}(X, \mathbb{Z}/\ell(2+n)) = E_n^{3+n} \supseteq E_{n+1}^{3+n} \supseteq 0$$

which leads to the exact sequence

$$0 \rightarrow E_\infty^{n+1,2} \longrightarrow H^{3+n}(X, \mathbb{Z}/\ell(2+n)) \longrightarrow E_\infty^{n,3} \longrightarrow 0$$

Since $E_2^{p,q} = 0$ for all $p \geq n+2$ or $q \geq 4$, we see that

$$E_2^{n,3} = E_3^{n,3} = \dots = E_\infty^{n,3}.$$

The same argument yields

$$E_3^{n+1,2} = E_4^{n+1,2} = \dots = E_\infty^{n+1,2}$$

and $E_3^{n+1,2} = \text{Coker } d_2^{n-1,3}$ where $d_2^{n-1,3}$ is the map

$$H^{n-1}(k, H^3(X', \mathbb{Z}/\ell(2+n))) \longrightarrow H^{n+1}(k, H^2(X', \mathbb{Z}/\ell(2+n))).$$

Hence, we obtain the exact sequence

$$(3.7) \quad 0 \rightarrow \text{Coker } d_2^{n-1,3} \longrightarrow H^{3+n}(X, \mathbb{Z}/\ell(2+n)) \longrightarrow H^n(k, H^3(X', \mathbb{Z}/\ell(2+n))) \longrightarrow 0$$

Combining duality (2.2) for k and duality (3.2), we deduce that the group $H^0(k, H^1(X', \mathbb{Z}/\ell))$ is dual to the group $H^{n+1}(k, H^2(X', \mathbb{Z}/\ell(2+n)))$ and the group $H^2(k, H^0(X', \mathbb{Z}/\ell))$ is dual to the group $H^{n-1}(k, H^3(X', \mathbb{Z}/\ell(2+n)))$. On the other hand, we have the commutative diagram

$$(3.8) \quad \begin{array}{ccccccc} H^{n-1}(k, H^3(X', \mathbb{Z}/\ell(2+n))) & \times & H^2(k, H^0(X', \mathbb{Z}/\ell)) & \longrightarrow & H^2(k, \mathbb{Z}/\ell(1)) & \xrightarrow{\sim} & \mathbb{Z}/\ell \\ & & \uparrow & & \parallel & & \parallel \\ H^{n+1}(k, H^2(X', \mathbb{Z}/\ell(2+n))) & \times & H^0(k, H^1(X', \mathbb{Z}/\ell)) & \longrightarrow & H^2(k, \mathbb{Z}/\ell(1)) & \xrightarrow{\sim} & \mathbb{Z}/\ell \end{array}$$

given by the cup products and the spectral sequence (3.3), using the same argument as ([1], diagram 46). We infer that $\text{Coker } d_2^{n-1,3}$ is the dual of $\text{Ker}' d_2^{0,1}$ where $d_2^{0,1}$ is the boundary map for the spectral sequence ((3.4), j=0)

$$(3.9) \quad {}'E_2^{p,q} = H^p(k, H^q(X', \mathbb{Z}/\ell)) \implies H^{p+q}(X, \mathbb{Z}/\ell)$$

Similarly, the group $H^n(k, H^3(X', \mathbb{Z}/\ell(2+n)))$ is dual to the group $H^1(k, H^0(X', \mathbb{Z}/\ell))$. The required duality is deduced from the following commutative diagram

$$\begin{array}{ccccccc} 0 \rightarrow \text{Coker } d_2^{n-1,3} & \longrightarrow & H^{3+n}(X, \mathbb{Z}/\ell(2+n)) & \longrightarrow & H^n(k, H^3(X', \mathbb{Z}/\ell(2+n))) & \longrightarrow & 0 \\ & \downarrow \wr & & \downarrow \wr & & \downarrow \wr & \\ 0 \rightarrow (\text{Ker}' d_2^{0,1})^\vee & \longrightarrow & (H^1(X, \mathbb{Z}/\ell))^\vee & \longrightarrow & (H^1(k, H^0(X', \mathbb{Z}/\ell)))^\vee & \longrightarrow & 0 \end{array}$$

where the upper exact sequence is (3.7) and the bottom exact sequence is the dual of the well-known exact sequence

$$0 \rightarrow {}'E_2^{1,0} \longrightarrow H^1(X, \mathbb{Z}/\ell) \longrightarrow \text{Ker}' d_2^{0,1} \longrightarrow 0$$

deduced from the spectral sequence (3.9) and where $(M)^\vee$ denotes the dual $\text{Hom}(M, \mathbb{Z}/\ell)$ for any $\mathbb{Z}/\ell$ -module M .

Finally, to obtain the last part of the theorem, we remark that the commutativity of the diagram (3.6) is obtained by via a same argument (projection formula ([9], VI 6.5) and compatibility of traces ([9], VI 11.1)) as [1] to establish the commutative diagram in the proof of assertion ii) at page 791. $\square$

Corollary 3.2

With the same notations as above, the following commutative diagram

$$(3.10) \quad \begin{array}{ccc} H^{n+1}(k(v), \mathbb{Z}/\ell(n+1)) & \xrightarrow{i_*} & H^{3+n}(X, \mathbb{Z}/\ell(2+n)) \\ \downarrow & & \downarrow \\ (H^1(k(v), \mathbb{Z}/\ell))^\vee & \xrightarrow{(i^*)^\wedge} & (H^1(X, \mathbb{Z}/\ell))^\vee \end{array}$$

holds.

Proof. This is a consequence of diagram (3.6). $\square$

The duality (3.5) will be completed (section 5) to a general pairing by replacing $H^1(X, \mathbb{Z}/\ell)$ by $H^i(X, \mathbb{Z}/\ell)$; for $0 \leq i \leq 5$ in the case $n = 1$.

4. THE BLOCH-OGUS COMPLEX

In this section, we investigate the study of the Bloch-Ogus complex associated to the ring A considered previously. So, let A be a 2-dimensional normal complete local ring whose residue field is an n -dimensional local field. Next, we define a group which appears in the homologies of the associated Bloch-Ogus complex of A .

Definition 4.1

Let Z be a Noetherian scheme. A finite etale covering $f : W \rightarrow Z$ is called a *c.s covering* if for any closed point z of Z , $z \times_Z W$ is isomorphic to a finite scheme-theoretic sum of copies of z . We denote $\pi_1^{c.s}(Z)$ the quotient group of $\pi_1^{ab}(Z)$ which classifies abelian c.s coverings of Z .

As above, let $X = \text{Spec} A \setminus \{x\}$. The group $\pi_1^{c.s}(X)/\ell$ is the dual of the kernel of the map

$$(4.1) \quad H^1(X, \mathbb{Z}/\ell) \longrightarrow \prod_{v \in P} H^1(k(v), \mathbb{Z}/\ell)$$

([12], section 2, definition and sentence just below). Now, we are able to calculate the homologies of the Bloch-Ogus complex associated to the ring A .

Theorem 4.2

For all ℓ prime to the characteristic of A , the following sequence is exact.

$$(4.2) \quad 0 \longrightarrow \pi_1^{c.s}(X)/\ell \longrightarrow H^{n+3}(K, \mathbb{Z}/\ell(n+2)) \longrightarrow \bigoplus_{v \in P} H^{n+2}(k(v), \mathbb{Z}/\ell(n+1)) \longrightarrow \mathbb{Z}/\ell \longrightarrow 0$$

Proof. Consider the localisation sequence on $X = \text{Spec} A \setminus \{x\}$

$$\dots \rightarrow H^i(X, \mathbb{Z}/\ell(n+2)) \rightarrow H^i(K, \mathbb{Z}/\ell(n+2)) \rightarrow \bigoplus_{v \in P} H_v^{i+1}(X, \mathbb{Z}/\ell(n+2)) \rightarrow \dots$$

Firstly, for any $v \in P$, we have the isomorphisms

$$H_v^i(X, \mathbb{Z}/\ell(2+n)) \simeq H_v^i(\text{Spec} A_v, \mathbb{Z}/\ell(2+n))$$

by excision. Secondly, we can apply the purity theorem of Fujiwara-Gabber (Introduction) for $Z = v, T = \text{Spec} A_v$ and we find the isomorphisms

$$H_v^{3+n}(\text{Spec} A_v, \mathbb{Z}/\ell(2+n)) \simeq H^{1+n}(k(v), \mathbb{Z}/\ell(1+n))$$

and

$$H_v^{4+n}(\text{Spec} A_v, \mathbb{Z}/\ell(2+n)) \simeq H^{2+n}(k(v), \mathbb{Z}/\ell(1+n))$$

which lead to the isomorphisms

$$H_v^{3+n}(X, \mathbb{Z}/\ell(2+n)) \simeq H^{1+n}(k(v), \mathbb{Z}/\ell(1+n))$$

and

$$H_v^{4+n}(X, \mathbb{Z}/\ell(2+n)) \simeq H^{2+n}(k(v), \mathbb{Z}/\ell(1+n)).$$

Hence we derive the exact sequence

$$\begin{aligned} \bigoplus_{v \in P} H^{1+n}(k(v), \mathbb{Z}/\ell(1+n)) &\xrightarrow{g} H^{3+n}(X, \mathbb{Z}/\ell(2+n)) \longrightarrow H^{3+n}(K, \mathbb{Z}/\ell(2+n)) \\ &\longrightarrow \bigoplus_{v \in P} H^{2+n}(k(v), \mathbb{Z}/\ell(1+n)) \longrightarrow H^{4+n}(X, \mathbb{Z}/\ell(2+n)) \longrightarrow 0 \end{aligned}$$

The last zero on the right is a consequence of the vanishing of the group

$H^{4+n}(K, \mathbb{Z}/\ell(2+n))$. Indeed, A is finite over $O_L[[T]]$ for some complete discrete valuation field L having the same residue field with A [10, §31]. By Serre [13, chapI, Prop 14], $cd_\ell(A) \leq cd_\ell(O_L[[T]])$ and by Gabber [5], $cd_\ell(O_L[[T]]) = n + 3$ using the fact that $cd_\ell(k) = n + 1$.

Now, by the right square of the diagram (3.6), the Gysin map

$$\bigoplus_{v \in P} H^{2+n}(k(v), \mathbb{Z}/\ell(1+n)) \longrightarrow H^{4+n}(X, \mathbb{Z}/\ell(2+n))$$

can be replaced by the map $\bigoplus_{v \in P} H^{2+n}(k(v), \mathbb{Z}/\ell(1+n)) \longrightarrow \mathbb{Z}/\ell$ after composing with the trace isomorphism $H^{4+n}(X, \mathbb{Z}/\ell(2+n)) \simeq \mathbb{Z}/\ell$ (3.4). So, we obtain the exact sequence

$$0 \longrightarrow \text{Coker } g \longrightarrow H^{3+n}(K, \mathbb{Z}/\ell(2+n)) \longrightarrow \bigoplus_{v \in P} H^{2+n}(k(v), \mathbb{Z}/\ell(1+n)) \longrightarrow \mathbb{Z}/\ell \longrightarrow 0$$

Finally, in view of the commutative diagram (3.10), we deduce that $\text{Coker } g$ equals to the group $\pi_1^{c,s}(X)/\ell$ taking in account (4.1). $\square$

Next, we assume further that A is regular. We will prove that the group $\pi_1^{c,s}(X)/\ell$ vanishes.

Theorem 4.3

Let $A = F_p((t_1))((t_2)) \dots ((t_n))[[X, Y]]$ of fraction field K and assum Kato conjecture (section 2), then the following Hasse principle complex for K

$$(4.3) \quad 0 \longrightarrow H^{3+n}(K, \mathbb{Z}/\ell(2+n)) \longrightarrow \bigoplus_{v \in P} H^{2+n}(k(v), \mathbb{Z}/\ell(1+n)) \longrightarrow \mathbb{Z}/\ell \longrightarrow 0$$

is exact.

Proof. Keeping in mind (4.2), it remains to prove the injectivity of the map

$$\Psi_K : H^{3+n}(K, \mathbb{Z}/\ell(2+n)) \longrightarrow \bigoplus_{v \in P} H^{2+n}(k(v), \mathbb{Z}/\ell(1+n)).$$

Let q be an integer and consider the sheaf $\mathcal{H}^q(\mathbb{Z}/\ell(n+2))$ on $\text{Spec} A$, the Zariskien sheaf associated to the presheaf $U \longrightarrow H^q(U, \mathbb{Z}/\ell(n+2))$. As a consequence of a recent work of Panin [11], we conclude that the cohomology of this sheaf is calculated as the homology of the Bloch-Ogus complex, that is :

$$H^q(K, \mathbb{Z}/\ell(n+2)) \longrightarrow \bigoplus_{v \in P} H^{q-1}(k(v), \mathbb{Z}/\ell(n+1)) \longrightarrow H^{q-1}(k(x), \mathbb{Z}/\ell(n)).$$

So the group $\text{Ker } \Psi_K$ is identified with the group $H^0((\text{Spec} A)_{\text{zar}}, \mathcal{H}^{n+3}(\mathbb{Z}/\ell(n+2)))$.

On the other hand, the Bloch-Ogus spectral sequence

$$H^p((\text{Spec} A)_{\text{zar}}, \mathcal{H}^q(\mathbb{Z}/\ell(n+2))) \Rightarrow H^{p+q}(\text{Spec} A, \mathbb{Z}/\ell(n+2))$$

gives the exact sequence

$$\begin{aligned} H^{n+3}(\text{Spec} A, \mathbb{Z}/\ell(n+2)) &\longrightarrow H^0((\text{Spec} A)_{\text{zar}}, \mathcal{H}^{n+3}(\mathbb{Z}/\ell(n+2))) \\ &\longrightarrow H^2((\text{Spec} A)_{\text{zar}}, \mathcal{H}^{n+2}(\mathbb{Z}/\ell(n+2))) \rightarrow H^{n+4}(\text{Spec} A, \mathbb{Z}/\ell(n+2)) \end{aligned}$$

Since the ring A is henselian, we obtain the isomorphism

$$H^i(\operatorname{Spec} A, \mathbb{Z}/\ell(n+2)) \simeq H^i(\operatorname{Spec} k, \mathbb{Z}/\ell(n+2)), i \geq 0.$$

But the groups $H^{n+3}(\operatorname{Spec} k, \mathbb{Z}/\ell(3))$ and $H^{n+4}(\operatorname{Spec} k, \mathbb{Z}/\ell(3))$ vanish because the cohomological dimension of k is $n+1$. Thus we get the isomorphism

$$H^0((\operatorname{Spec} A)_{\operatorname{zar}}, \mathcal{H}^{n+3}(\mathbb{Z}/\ell(n+2))) \xrightarrow{\sim} H^2((\operatorname{Spec} A)_{\operatorname{zar}}, \mathcal{H}^{n+2}(\mathbb{Z}/\ell(n+2)))$$

which means that $\operatorname{Ker} \Psi_K$ is isomorphic to the Cokernel of the map

$$\bigoplus_{v \in P} H^{n+1}(k(v), \mathbb{Z}/\ell(n+1)) \longrightarrow H^n(k(x), \mathbb{Z}/\ell(n))$$

So, we must prove the surjectivity of this last map. Indeed, the Gersten-Quillen complex ([11], Theorem A)

$$K_{n+2}(A)/\ell \longrightarrow K_{n+2}(K)/\ell \longrightarrow \bigoplus_{v \in P} K_{n+1}k(v)/\ell \longrightarrow K_n k(x)/\ell \longrightarrow 0$$

is exact. On the other hand, we have the following commutative diagram

$$\begin{array}{ccccccc} K_{n+2}K/\ell & \longrightarrow & \bigoplus_{v \in P} K_{n+1}k(v)/\ell & \longrightarrow & K_n k(x)/\ell & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow \wr & & \\ H^{n+2}(K, \mathbb{Z}/\ell(n+2)) & \longrightarrow & \bigoplus_{v \in P} H^{n+1}(k(v), \mathbb{Z}/\ell(n+1)) & \longrightarrow & H^n(k(x), \mathbb{Z}/\ell(n)) & & \end{array}$$

where the right vertical isomorphism comes from Kato conjecture. This yields that the map

$$\bigoplus_{v \in P} H^{n+1}(k(v), \mathbb{Z}/\ell(n+1)) \longrightarrow H^n(k(x), \mathbb{Z}/\ell(n))$$

is surjective and we are done. $\square$

Remark 4.4

1) The case $n = 0$, implies the following exact sequence

$$0 \longrightarrow H^3(K, \mathbb{Z}/\ell(2)) \longrightarrow \bigoplus_{v \in P} H^2(k(v), \mathbb{Z}/\ell(1)) \longrightarrow \mathbb{Z}/\ell \longrightarrow 0$$

already obtained by Saito [12].

2) The case $n = 1$ leads to the following exact sequence

$$0 \longrightarrow H^4(K, \mathbb{Z}/\ell(3)) \longrightarrow \bigoplus_{v \in P} H^3(k(v), \mathbb{Z}/\ell(2)) \longrightarrow \mathbb{Z}/\ell \longrightarrow 0$$

which is considered in [1].

3) The case $n = 2$. Let $A = \mathbb{F}_p((t))((u))[[X, Y]]$ of fraction field K . Then the following Hasse principle complex for K

$$0 \longrightarrow H^5(K, \mathbb{Z}/\ell(4)) \longrightarrow \bigoplus_{v \in P} H^4(k(v), \mathbb{Z}/\ell(3)) \longrightarrow \mathbb{Z}/\ell \longrightarrow 0$$

is exact.

4) The case $n = 0$ has been used by Saito to study the class field theory of curves over one dimensional local field. Recently, Yoshida [16] provided an alternative approach which includes the equal characteristic case. In a forthcoming paper I use the case $n = 1$ to investigate the study of class field theory of curves over 2- dimensional local field.

Let A denote a 2-dimensional complete normal local ring of positive characteristic whose residue field is one-dimensional local field. The aim of this section is to complete the duality (3.5) for $i \geq 1$. We prove the following.

Theorem 5.1

For all ℓ prime to $\text{char}(A)$, the isomorphism

$$(5.1) \quad H^5(X, \mathbb{Z}/\ell(3)) \simeq \mathbb{Z}/\ell$$

and the perfect pairing

$$(5.2) \quad H^i(X, \mathbb{Z}/\ell) \times H^{5-i}(X, \mathbb{Z}/\ell(3)) \longrightarrow H^5(X, \mathbb{Z}/\ell(3)) \simeq \mathbb{Z}/\ell$$

hold for all $i \in \{0, \dots, 5\}$.

Proof. The first isomorphism is given by (3.4). Next. We proceed to the second part of the theorem. As in the proof of Theorem 3.1, we consider the strict henselization A^{sh} of A (with respect to the separably closed extension k_s of k) at x . If x' is the maximal ideal of A^{sh} , we recall that we denote $X' = \text{Spec} A^{sh} \setminus \{x'\}$ and we consider the spectral sequence (3.3). The filtration of the group $H^i(X, \mathbb{Z}/\ell(3))$ is

$$H^i(X, \mathbb{Z}/\ell(3)) = E_0^i \supseteq E_1^i \supseteq E_2^i \supseteq 0$$

where the quotients are given by

$$E_0^i/E_1^i \simeq E_\infty^{0,i} \simeq \ker d_2^{0,i}$$

$$E_1^i/E_2^i \simeq E_\infty^{1,i-1} \simeq E_2^{1,i-1}, \text{ and}$$

$$E_2^i \simeq E_\infty^{2,i-2} \simeq \text{Coker } d_2^{0,i-1}$$

The same computation is true for the group $H^{5-i}(X, \mathbb{Z}/\ell)$ and the filtration

$$H^{5-i}(X, \mathbb{Z}/\ell) = {}'E_0^{5-i} \supseteq {}'E_1^{5-i} \supseteq {}'E_2^{5-i} \supseteq 0$$

by considering the spectral sequence ((3.4), $j=0$).

Now, combining duality (2.2) and duality (3.2) we observe that the group $E_2^{0,j}$ is dual to the group $'E_2^{2,3-j}$ and the group $E_2^{1,j}$ is dual to the group $'E_2^{1,3-j}$ for all $0 \leq j \leq 3$. On the other hand, we have the commutative diagram

$$\begin{array}{ccccccc} H^0(k, H^i(X', \mathbb{Z}/\ell(3))) & \times & H^2(k, H^{3-i}(X', \mathbb{Z}/\ell)) & \longrightarrow & H^2(k, \mathbb{Z}/\ell(1)) & \xrightarrow{\sim} & \mathbb{Z}/\ell \\ \downarrow & & \uparrow & & \parallel & & \parallel \\ H^2(k, H^{i-1}(X', \mathbb{Z}/\ell(3))) & \times & H^0(k, H^{4-i}(X', \mathbb{Z}/\ell)) & \longrightarrow & H^2(k, \mathbb{Z}/\ell(1)) & \xrightarrow{\sim} & \mathbb{Z}/\ell \end{array}$$

given by the cup products and the spectral sequence (3.4), using the same argument as ([1], diagram 46). We infer that $\text{Coker } d_2^{0,i-1}$ is the dual of $\text{Ker}' d_2^{0,5-i}$ and $\text{Ker } d_2^{0,i}$ is the dual of $\text{Coker}' d_2^{0,2d-i}$ where $'d_2^{p,q}$ is the boundary map for the spectral sequence ((3.4), $j=0$)

$$'E_2^{p,q} = H^p(k, H^q(X', \mathbb{Z}/\ell)) \implies H^{p+q}(X, \mathbb{Z}/\ell).$$

This is illustrated by the following diagram:

$$\begin{array}{ccccccc}
(5.3) & H^i X, \mathbb{Z}/\ell(3)) = E_0^i & & & & & 0 \\
& \downarrow & \} \text{Ker} d_2^{0,i} & \longleftrightarrow & \text{Co ker}' d_2^{0,24-i} \{ & & \downarrow \\
& E_1^i & & & & & 'E_2^{5-i} \\
& \downarrow & \} E_2^{1,i-1} & \longleftrightarrow & 'E_2^{1,24-i} \{ & & \downarrow \\
& E_2^i & & & & & 'E_1^{5-i} \\
& \downarrow & \} \text{Co ker} d_2^{0,i-1} & \longleftrightarrow & \text{Ker}' d_2^{0,5-i} \{ & & \downarrow \\
& 0 & & & & & 'E_0^{5-i} = H^{5-i}(X, \mathbb{Z}/\ell)
\end{array}$$

where each pair of groups which are combined by $\longleftrightarrow$ consists of a group and its dual group.

We begin by calculate the dual group of E_1^i , using the following commutative diagram

$$\begin{array}{ccccccc}
0 \rightarrow & E_2^i & \rightarrow & E_1^i & \rightarrow & E_1^i/E_2^i & \rightarrow 0 \\
& \downarrow \wr & & \downarrow & & \downarrow \wr & \\
0 \rightarrow & ('E_0^{2d+1-i}/'E_1^{2d+1-i})^\vee & \rightarrow & ('E_0^{2d+1-i}/'E_2^{2d+1-i})^\vee & \rightarrow & ('E_1^{2d+1-i}/'E_2^{2d+1-i})^\vee & \rightarrow 0
\end{array}$$

where $(M)^\vee$ denotes the dual $\text{Hom}(M, \mathbb{Z}/\ell)$ for any $\mathbb{Z}/\ell$ -module M and where the left and right vertical isomorphisms are explained by the previous diagram. This yields that

$$(5.4) \quad E_1^i \simeq ('E_0^{5-i}/'E_2^{5-i})^\vee$$

Finally, the required duality between $H^i X, \mathbb{Z}/\ell(d+1)) = E_0^i$ and $'E_0^{5-i} = H^{5-i}(X, \mathbb{Z}/\ell)$ follows from the following commutative diagram

$$\begin{array}{ccccccc}
0 \rightarrow & E_1^i & \rightarrow & E_0^i & \rightarrow & E_0^i/E_1^i & \rightarrow 0 \\
& \downarrow \wr & & \downarrow \wr & & \downarrow \wr & \\
0 \rightarrow & ('E_0^{5-i}/'E_2^{5-i})^\vee & \rightarrow & ('E_0^{5-i})^\vee & \rightarrow & ('E_2^{5-i})^\vee & \rightarrow 0
\end{array}$$

where the right vertical isomorphism is given by (5.3) and the left vertical isomorphism is the isomorphism (5.4). $\square$

REFERENCES

- [1] Colliot-Thélène, J.L., Sansuc, J.J., Soulé, C. Torsion dans le groupe de Chow de codimension deux. Duke Math. Journal vol. 50 No.3 pp763-801 (1983)
- [2] Draouil, B. Cohomological Hasse principle for the ring $\mathbb{F}_p((t))[[X, Y]]$, Bull. Belg. Math. Soc. Simon Stevin 11, no. 2 (2004), 181–190
- [3] Draouil, B., Douai, J.C. Sur l'arithmétique des anneaux locaux de dimension 2 et 3, Journal of Algebra 213 (1999), 499-512.
- [4] Fujiwara, K. Theory of Tubular Neighborhood in Etale Topology Duke Math. J. 80 (1995), 15-57.
- [5] Gabber, O. Lecture at IHES, on March 1981.
- [6] Kato, K. A Hasse principle for two-dimensional global fields, J. reine angew. Math. 366 (1986), 143-183.
- [7] Kato, K. A generalisation of local class field theory by using K-theory II, J. Fac. Sci. Univ. Tokyo, 27 (1980), 603-683.
- [8] Matsumi, K. Thesis, Arithmetic of three-dimensional complete regular local rings of positive characteristics T_0^2 hoku University, Japan 1999.
- [9] Milne, J.S. Etale Cohomology, Princeton University Press, Princeton 1980.
- [10] Nagata, M. Local rings, Tracts in Mathematics Number 13, Intersciences Publishers. New York 1962
- [11] Panin, I. The equi-characteristic case of the Gersten conjecture, preprint (2000), available on the K-theory server.

- [12] Saito, S. Class field Theory for two-dimensional local rings Galois groups and their representations, Kinokuniya-North Holland Amsterdam, vol 12 (1987), 343-373
- [13] Serre,J.P. Cohomologie Galoisienne ,L.N.M 5 Berlin-Heidelberg-New York 1965.
- [14] SGA 4 Théorie des Topos et Cohomologie étale des Schémas. Lecture Notes in Math. vol.305, Springer-Verlag Berlin.Heidelberg.New York.
- [15] SGA 5 Cohomologie l-adique et Fonctions L . Lecture Notes in Math. vol.589, Springer-Verlag Berlin.Heidelberg.New York
- [16] Yoshida. Finiteness theorems in the class field theory of varieties over local fields, Journal of Number Theory 101 (2003) 138-150

DÉPARTEMENT DE MATHÉMATIQUES, FACULTÉ DES SCIENCES DE BIZERTE, 7021 JARZOUNA

TUNISIA

E-mail address: Belgacem.Draouil@fsb.rnu.tn